Prevalence of Bacterial Urinary Tract Infections and Associated Factors among Patients Attending Hospitals

SYNOPSIS
SUBMITED TO

Jayoti Vidyapeeth Women's University

In partial fulfillment of the requirements for the degree of

DOCTOR OF PHILOSOPHY
IN

Microbiology
By

JV'n Rupal Sengar

UNDER THE SUPERVISION OF

GUIDE- JV'n Dr. Parita
Associate Professor
Jayoti Vidyaurineth Women's University

CO-GUIDE- DrAnil Kumar
Assistant professor,
SBS University,dehradun

Jayoti Vidyaurineth Women's University, Jaipur (Rajasthan)
June, 2019

INTRODUCTION

Kidney produces the waste product urine.. It is a reasonable, straightforward liquid that ordinarily has a golden shading. Synthetically, **urine** is essentially a watery arrangement of salt and substances called urea and uric corrosive.In any case, microbes can enter the urethra and cause a disease (UTI). As a result of its positive synthetic creation human **urine** can bolster development of a few distinct strains of microscopic organisms (Srikanta Chowdhury and Ramendu Parial, 2014). The **urine** culture is utilized to analyze a urinary tract disease (UTI) and to recognize the microscopic organisms or yeast causing the contamination. It might be done related to defenselessness testing to figure out which anti-toxins will restrain the development of the organism causing the disease.

UTI alludes to the presence of microbial pathogens in the urinary tract and is characterized as the development of a solitary pathogen of >105 settlement framing units per milliliter from appropriately gathered mid-stream **urine** examples (Zahra Tayebi et.al, 2016). Urinary tract contamination (UTI) is the disease of any piece of body's excretory framework which incorporates two kidneys, two ureters, a bladder, and a urethra. Urinary tract disease (UTI) is the second most basic medical clinic obtained and network gained contaminations after respiratory tract contaminations with an expected yearly worldwide rate of at any rate 250 million cases and almost 10% of individuals will encounter an UTI during their lifetime (Md. Abdullah Yusuf et.al, 2015, B Khatri et.al, 2012, Inam Ullah Khan et.al, 2014, Arunima Moue et.al, 2015, Nayareen Akhtar et.al, 2016). Nosocomial contaminations are significant general medical issues in creating nations. Nosocomial or clinic procured diseases are generally characterized as contaminations that are distinguished at any rate 48-72 hours following admission to medical clinic and medicinal services office. Emergency clinic procured UTIs are a genuine risk particularly for insusceptible traded off patients as it can cost a critical monetary weight to the medical clinic the executives (Inam Ullah Khan et.al, 2014). Bacterial specialists are in charge of UTI that can be gone from gentle irritative voiding to bacteremia, sepsis and demise (Zahra Tayebi et.al, 2016).

Expansion of microorganisms in the urinary tract is the reason for urinary tract contamination. The clinical signs of UTI rely upon the piece of the urinary tract included, the etiologic living beings(Saima Mollick et.al, 2016). Gram negative microscopic organisms are common cause of uti(Md. Abdullah Yusuf et.al, 2015). They are cosmopolitan in nature and ready to start, set up and cause different sorts of contaminations that is the reason these living beings have an extraordinary

clinical significance. Creatures that reason UTI are those from the ordinary vaginal, perineal, and fecal vegetation. Proteus mirabilis, Pseudomonas aeruginosa, Acinetobacter, Serratia, Morganella margani, coagulase negative staphylococci and Streptococcus agalactiae (Arunima Moue et.al, 2015). Escherichia coli is the most predominant causative living being of UTI and is exclusively in charge of over 80% of these contaminations (Niranjan V. and Malini A., 2014, Inam Ullah Khan et.al, 2014, Nayareen Akhtar et.al, 2016). The strains of E. coli causing UTI have been named as Uropathogenic E. coli (UPEC). In UPEC strains, harmfulness variables incorporate the capacity to hold fast to uroepithelial cells, some O and K antigens, and protection from phagocytosis and to the bactericidal activity of human serum (Mucheye Gizachew et.al, 2013). Expanded Spectrum Beta-Lactamase (ESBL), Metallo Beta-Lactamase (MBL) and Amp-C interceded beta-lactamases are a portion of the chemicals created by (Inam Ullah Khan et.al, 2014).

This issue traverses all age gatherings, starting from the neonates to the geriatric age gathering (Dnyaneshwari Puroshottam Ghadage et.al, 2016). The predominance is expanded by a few elements. Poor financial status is accounted for to be a noteworthy hazard factor with needy patients having a fivefold expanded hazard. Other hazard variables incorporate expanded age, high equality, poor perineal cleanliness, history of repetitive UTI, diabetes mellitus, neurogenic bladder maintenance, anatomic or useful urinary tract variation from the norm, and expanded recurrence of sexual movement (Nayareen Akhtar et.al, 2016). UTIs are more typical in ladies and young ladies than in men because of anatomical and physiological reason (Md. Abdullah Yusuf et.al, 2015, Inam Ullah Khan et.al, 2014). This enables microbes from the digestive organs to come into contact all the more effectively with the urethra. Pregnancy additionally makes the ladies progressively vulnerable to the disease. Men likewise have an antibacterial substance in their prostate organ that brings down their hazard. It has been seen that about 20.0% of the ladies encountered a solitary scene of UTI during their lifetime, and 3.0% of ladies had more than one scene of UTI every year (Md. Abdullah Yusuf et.al, 2015, B Khatri et.al, 2012). Urinary tract disease (UTI) is likewise of incredible clinical significance attributable to impressively high grimness and death rates among kids (Shahla Afsharpaiman et.al, 2012). The diseases might be symptomatic or asymptomatic, and either kind of contamination can result in genuine sequelae whenever left untreated. The most widely recognized indications for which most patients look for treatment are dysuria and recurrence (B Khatri et.al, 2012). Pediatric UTIs are related with high horribleness and long haul complexities like renal scarring, hypertension and endless renal disappointment. Determination is troublesome in the neonatal period on the grounds that the signs and manifestations are non-explicit in this age gathering

(Dnyaneshwari Puroshottam Ghadage et.al, 2016).

Antibacterial Resistance-

With the approach of novel anti-microbial is performed.In practically all instances of UTI, exact antimicrobial treatment starts before the lab aftereffects of **urine** culture are accessible; accordingly anti-microbial obstruction may increment in uropathogens because of regular utilization of anti-microbials (Ruchita Mahajan et.al, 2014). Example of anti-microbial affectability may change even over brief periods furthermore, from spot to put, and intermittent assessment of anti-microbial affectability is expected to refresh this data (Kiran K Mokta et.al, 2015). Protection from anti-microbials is exceedingly pervasive in bacterial disengages around the world, especially in creating nations (Manjula Mehta et.al, 2012Some populace based have demonstrated that protection from generally utilized anti-microbials, for example, trimethoprim-sulfamethoxazole, has multiplied in uropathogens. Comparable investigations on youthful Iranian kids with UTI have likewise uncovered uniquely high protection from generally utilized anti-toxins (Shahla Afsharpaiman et.al, 2012). These opposition properties are effectively moved between microscopic organisms of various genera through plasmids and different methods. To guarantee suitable treatment, learning of the creatures that reason UTI and their anti-microbial defenselessness is obligatory (Nayareen Akhtar et.al, 2016). Treatment depends on data decided from the antimicrobial opposition example of the urinary pathogens. In any case, as a result of the developing and proceeding with anti-infection opposition marvel, normal checking of obstruction examples is important to improve rules for observational anti-infection treatment (Nayareen Akhtar et.al, 2016). To be fruitful, the observational treatment gave must be guided by clinical confirmations just as the security profile and cost-viability of the medication, and cling to antimicrobial stewardship (Kiran K Mokta et.al, 2015).

Review of Literature

Taiwo SS and Aderounmu AOA, in 2006, recognized microbial pathogens related with bacteriuria and UTI in patients with inhabiting urethral catheters and decide their helplessness examples to normally utilized antimicrobial operators in their organization. They biochemically described unadulterated provinces of detached life forms and performed plate dissemination anti-toxin vulnerability on each bacterial disengage.Klebsiella spp were the commonest pathogen confined with 46 (36.6%), trailed by Pseudomonas spp 34 (27.0%), Escherichia coli 26 (20.6%),. They inferred that catheter-related UTI in their organization was brought about by multi-safe microbial pathogens which had happened subsequent on prophylactic anti-microbial treatment regulated after catheterization. Accentuation ought to be set on great catheter the board as opposed to the utilization of prophylaxis, to decrease the frequency of catheter related UTI.

Mansour Amin et.al, in 2009, considered on microscopic organisms confined from urinary tract diseases and decided their powerlessness to anti-infection agents. He found that roughly 1 out of 3 ladies would require antimicrobial treatment for a UTI before age 24, and 40% to half of ladies would have an UTI during their lifetime.

. Urine examples were refined for separation of the microbial operators of UTI. The segregated microscopic organisms were recognized utilizing biochemical tests. Circle dissemination defenselessness test was utilized to decide weakness of bacterial operators to anti-infection agents. In that review 553(8.7%) patients out of 7056 were demonstrated to be urine culture positive (68% females and 32% guys). The most detached bacterium was E. coli with recurrence rate of 59%. The commonness of UTIs in the referral clinic was 67.2%. The occurrence was higher in females with a pervasiveness rate of 54.3%, while in guys the recorded worth was 45.7%. Gram-negative secludes had a commonness of 74.7%, while gram-positive segregates had 25.30%. Different less-visit confines in total caused 16.1% of diseases. They performed anti-toxin affectability and opposition examination by the circle dissemination technique utilizing business anti-microbials plates. Powerlessness of the clinical detaches to anti-microbials generally utilized in the 750 bed referral emergency clinic was variable, contingent upon

species and medication being referred to. The mean weakness of the segregates recorded were E. coli (36.3%), K. pneumoniae (44.1%), P. mirabilis (47.8%), S. faecalis (51.3%), S. aureus (43.1 %), P. vulgaris (48.1%), P. stuartii (45.0%), S. epidermidis (49.7%), Al. faecalis (54.5%), S. saprophyticus (54.2%), P. aeruginosa (38.6%), S. marcescens (44.2%) and C. fruendii (40.4%). The mean affectability of the anti-toxins were ofloxacin (63.8%), gentamycin (13.26%), streptomycin (37.0%), colistin (49.2%), ampicillin (25.4%), nalidixic corrosive (45.5%), nitrofurantoin (55.4%), augmentin (64.9%), antibiotic medication (27.0%), cotrimoxazole (41.8%), pefloxacin (34.9%), chloramphenicol (50.9%), and

erythromycin (51.6%). The information got in this investigation featured the issues of bacterial predominance and opposition among uropathogenic microbes in Yola, Nigeria.

Shahla Afsharpaiman et.al, in 2012, completed a cross sectional examination on bacterial pathogens and obstruction designs in youngsters with network procured urinary tract contamination and found that UTI in kids had indicated elevated amounts of anti-microbial opposition in clinical settings. The investigation subjects were 114 kids (female subjects, 58.8%) with culture-demonstrated UTI sorted in the accompanying 3 age gatherings: neonates (< 28 days, n = 45), babies (28 days to 2 years, n = 41), and youngsters (> 2 years, n = 28). The most oftentimes refined pathogens included Escherichia coli (71.7%) and Enterobacter (28.9%). UTI brought about by Enterobacter was all the more ordinarily distinguished in neonates (60.6%) than in newborn children (21.2%) and youngsters (18.2%). Imipenem was the most dynamic operator against E. coli confines (helplessness, 97.3%), trailed by ciprofloxacin (90.4%) and amikacin .
Kemebradikumo Pondei1 et.al, in 2012, decide the normal causative operators of UTI and their antimicrobial defenselessness designs in a little however bustling private emergency clinic in Bayelsa State of Nigeria. Clean-get mid-stream urine tests were acquired from 116 investigation subjects.. Bacterial detaches were recognized morphologically and by standard biochemical tests. Anti-microbial vulnerability was tried utilizing the circle dispersion method of Kirby-Bauer. 36 tests out of 116 were culture positive trailed by Escherichia coli (36.1%). There was female prevalence of UTI and expanded frequency in the 21 to 30 years age gathering. Bacterial disconnects were delicate to gentamicin and amoxicillin-clavulanic corrosive, yet were impervious to nalidixic corrosive. They presumed that distinctions exist in the causative specialists of UTI and their antimicrobial affectability designs between social insurance offices. They suggest that every office ought to decide these files to direct their administration of uncomplicated UTI.

B Khatri, et.al, in 2012, depicted the etiological specialists of UTI and their commonness, and decided the antimicrobial weakness example of the bacterial pathogens segregated from urine culture. Midstream Urine tests from 1323 patients associated with UTI were examined by microscopy, and customary semi-quantitative culture method for the noteworthy development. Antimicrobial helplessness test was performed for the confines by Modified Kirby-Bauer plate dissemination strategy. The general predominance of UTI was observed to be 18.89%.

Aim-

To determine Occurrence of Bacterial Urinary System Infections and Associated Aspects among Patients Attending Hospital .

Objective-

1. To perform the isolation of bacteria .
2. To study the confirmation of Bacteria.
3. To perform the Biochemical test for the confirmation of bacteria.
4. To assess the Antibacterial Sensitivity Test.
5. To compare the bacterial sensitivity to antibiotics according to growth in the media.

MATERIAL AND METHODS

The present study is intended to carried out in Dept of Microbiology,MBS Hospital, Kota (Rajasthan) .This will include a minimum nuber of 100 pateints screened for bacterial disease of UTI in hospital attending patients of MBS Hospital.
All the informed pertaining to their disease will be recorded. The ethical clearance will be taken from IEC(Institutional ethical committee) before starting my work.
Inclusion criteria- The sample will be taken of 15 years above ,diabetics patients.
Exclusion criteria- Patient having age of below 15 years, pregnant women these patients will exclude form the study.
After taking informed consent from the patient the urine sample collection is done in sterile container. The patient will be guided to collect first voided midstream early morning urine sample in the sterile plastic container after cleaning of external genitalia. The mid portion of urine (30 ml) should be collected directly into the sterile leak proof container. Close the container immediately. The sample will be proceeded within 1 hr after collection for the further storage of sample to verify the results the preservative will be added. Preservatives for culture and antibiotic susceptibility testing are made to make the specimen in a fresh stage that can be achieved with refrigeration by the multiply of organisms this would result in wrong bacteria culture or bacterial overgrowth. Thing to consider must be given {in order to the formulation of the chemical |preservatives to achieve this target. There is certainly evidence to recommend that non-pH buffered boric acid might be harmful in order to certain organisms and this buffered boric acid preservatives can easily reduce the linked} with the preservative in the organisms. Preserved specimens may be stored with room temperature until typically the time of testing. Suitable labeling should be performed. the label needs to be located on the container but not on the lid. Typically the label will probably be properly affixed to the container.

The sample will be further processed for selected parameters:-

1.Isolation of bacteria- Isolation of bacteria will be done on media-

1.Cysteine Lactose Electrolyte (CLED) agar media
2. 5% Sheep Blood agar media
3.Nutrient Agar

2.Identification of bacteria-

For the bacterial identification staining technique will be applied over it. Grams staining will be the best method to check the result of samples.
Reference- The protocol of Microscope master will be followed in the study.

3. To perform the Biochemical test for the confirmation of bacteria.

Different Biochemical test will be done to check the bacteria. The confirmation of the bacteria can be done by by biochemical test. The protocol will be followed by A database (VetBact/VetBakt).

4. To assess the Antibacterial Sensitivity Test

The manual technique involves the Kirby Bauer or Disc Diffusion technique will be taken in this study.

The outcome of Research Work:-

1. The outcome of this research work is expected to be useful to the physician to predict the severity of decease as well to safe of patients who are on the way to progress the bacterial-induced UTI infection.
2. It can also give the effectiveness to multidrug therapy in future aspect.

REFERENCE

- Arunima Moue *et.al;* Prevalence of urinary tract infection in both outpatient department and in patient department at a medical college setting of Bangladesh. International Journal of Biosciences. 2015; 7(5); p. 146-152.
- B Khatri *et.al;* Etiology and antimicrobial susceptibility pattern of bacterial pathogens from urinary tract infection. Nepal Med Coll J 2012; 14(2): 129-132.
- Cunha MA *et.al;* Antibiotic resistance patterns of urinary tract infections in a northeastern Brazilian capital. Rev Inst Med Trop Sao Paulo. 2016; 58:2.
- Dnyaneshwari P. Ghadage *et.al;* Bacteriological Profile and Antibiogram of Urinary Tract Infections at a Tertiary Care Hospital. National Journal of Laboratory Medicine. 2016; 5(4): MO20-MO24
- El-Mahmood Muhammad Abubakar *et.al;* Antimicrobial susceptibility pattern of pathogenic bacteria causing urinary tract infections at the Specialist Hospital, Yola, Adamawa state, Nigeria. Journal of Clinical Medicine and Research. 2009; 1(1) pp. 001- 008
- Goli Angoti *et.al;* Bacteria Isolated from Urinary Tract Infection among Patients and Determination of the Antibiotic Susceptibility Patterns of the Gram Negative Bacteria in Iran. Novelty in Biomedicine. 2016; 1, 1-4.
- Gupta V *et.al;* Resistance Pattern in Uropathogens. Indian J Med Microbiol, 2002; 20: 96-8.
- Inam Ullah Khan *et.al;* (2014); Antimicrobial Susceptibility Pattern of Bacteria Isolated from Patients with Urinary Tract Infection; Journal of the College of Physicians and Surgeons Pakistan, 2014; Vol. 24 (11): 840-844.

- . M. Al-Jebouri and S. A. Mdish. Antibiotic Resistance Pattern of Bacteria Isolated from Patients of Urinary Tract Infections in Iraq. Open Journal of Urology. 2013, 3, 124- 131.
- Mahim Koshariya *et.al,* Prevalence of pathogens and their antimicrobial susceptibility in catheter associated urinary tract infection; IAIM, 2015; 2(4): 96-113.
- Manjula Mehta *et.al;* Prevalence and Antibiotic Susceptibility Pattern of Multi-Drug Resistant Escherichia coli Isolates from Urinary Tract Infection Patients. International Journal of Life Science and Pharma research. 2012; 2:4: 2250-0480.
- Md. Abdullah Yusuf *et.al* (2015); Antibiotic sensitivity pattern of gram negative uropathogenic bacilli at a private hospital in Dhaka city; Al Ameen J Med Sci; 8(3):189- 194.
- Mokta *et.al;* Bacterial Etiology and Antibiotic Susceptibility Pattern of Urinary Tract Infection. National Journal of Medical and Allied Sciences. 2015; 4(1)

- Mollick *et al;* Isolation and Characterization of Pathogens Responsible for Urinary Tract Infection in Bangladesh and Determination of their Antibiotic Susceptibility Pattern. Journal of Applied Pharmaceutical Science. 2016; 6 (04); 072-076.
- Mucheye Gizachew *et.al;* Escherichia coli isolated from patients suspected for urinary tract infections in Hawassa Referral Hospital, Southern Ethiopia: An institution based cross sectional study. E3 Journal of Microbiology. 2013; 1(1). pp. 009-015
- Nayareen Akhtar *et.al;* Antimicrobial Sensitivity Pattern of Escherichia coli Causing Urinary Tract Infection in Bangladeshi Patients. American Journal of Microbiological Research. 2016; 4(4); 122-125
- Niranjan V. & Malini A. Antimicrobial resistance pattern in Escherichia coli causing urinary tract infection among inpatients. Indian J Med Res 139, June 2014, pp 945-948.
- Shahla Afsharpaiman *et.al;* Bacterial Pathogens and Resistance Patterns in Children With Community-Acquired Urinary Tract Infection: A Cross Sectional Study. J Compr Ped.2012;3(1):16-20.
- Srikanta Chowdhury and Ramendu Parial, Antibiotic Susceptibility Patterns of Bacteria among Urinary Tract Infection Patients in Chittagong, Bangladesh. SMU Medical Journal. 2015; 2(1); 2349-1604
- Zahra Tayebi *et.al;* Frequency and antibiotic resistance pattern in gram positive uropathogenes isolated from hospitalized patients with urinary tract infection in Tehran, Iran. Journal of Genes, Microbes and Immunity. 2014 (2014); 1-9.
- https://vlab.amrita.edu/?sub=3&brch=73&sim=208&cnt=2(Gramstaining)